



Department of Biotechnology
Ministry of Science and Technology
Government of India
DBT



National Institute of
Advanced Industrial Science
and Technology
AIST

DBT - AIST International Laboratory for Advanced Biomedicine

DAILAB

Classroom for Advanced & Frontier Education CAFE

DAILAB-CAFE


Series - 07

Date and Time - February 9, 2015 (16:00~17:00) Venue - Central 4 (5F) Room 5105
Speaker – Masami Kojima
Affiliation – National Institute of Advanced Industrial Science and Technology (AIST), Japan
E-mail: m-kojima@aist.go.jp



Title – Single nucleotide polymorphism and proteolytic processing multiply the biological actions of growth factor in brain: A new insight into the growth factor theory

Most growth factors are initially synthesized as precursors and subsequently processed into their mature form by proteolytic cleavage, resulting in simultaneous removal of a pro-peptide. However, as compared to that of mature form, the biological role of the pro-peptide is poorly understood. Here, we investigated the biological role of the pro-peptide of brain-derived neurotrophic factor (BDNF) and first showed that the pro-peptide is expressed and secreted in hippocampal tissues and cultures, respectively. Interestingly, we found that the BDNF pro-peptide directly facilitates hippocampal long-term depression (LTD), requiring the activation of GluN2B-containing N-methyl-D-aspartate (NMDA) receptors and the pan-neurotrophin receptor p75^{NTR}. The BDNF pro-peptide also enhances NMDA-induced alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionic acid receptor endocytosis, a mechanism crucial for LTD expression. Thus, the BDNF pro-peptide is involved in synaptic plasticity that regulates a mechanism responsible for promoting LTD. The well-known BDNF polymorphism Val66Met affects human memory function. Here, the BDNF pro-peptide with Met mutation completely inhibits hippocampal LTD. These findings demonstrate functional roles for the BDNF pro-peptide and a naturally occurring human BDNF polymorphism in hippocampal synaptic depression.





Dear Dr Kojima

Thank you !

for being CAFÉ 07 Speaker

We all enjoyed your talk and appreciate your

efforts !

DAILAB-CAFE

Department of Biotechnology
Ministry of Science and Technology
Government of India
DBT

National Institute of
Advanced Industrial Science
and Technology
AIST

DBT - AIST International Laboratory for Advanced Biomedicine

DAILAB

Classroom for Advanced & Frontier Education CAFE

DAILAB-CAFE

Series - 07
 Date and Time - February 9, 2015 (16:00~17:00) Venue - Central 4 (5F) Room 5105
 Speaker - Masami Kojima
 Affiliation - National Institute of Advanced Industrial Science and Technology (AIST), Japan
 E-mail: m-kojima@aist.go.jp

Title - Single nucleotide polymorphism and proteolytic processing multiply the biological actions of growth factor in brain: A new insight into the growth factor theory

Most growth factors are initially synthesized as precursors and subsequently processed into their mature form by proteolytic cleavage, resulting in simultaneous removal of a pro-peptide. However, as compared to that of mature form, the biological role of the pro-peptide is poorly understood. Here, we investigated the biological role of the pro-peptide of brain-derived neurotrophic factor (BDNF) and first showed that the pro-peptide is expressed and secreted in hippocampal tissues and cultures, respectively. Interestingly, we found that the BDNF pro-peptide directly facilitates hippocampal long-term depression (LTD), requiring the activation of GluN2B-containing N-methyl-D-aspartate (NMDA) receptors and the pan-neurotrophin receptor p75^{NTR}. The BDNF pro-peptide also enhances NMDA-induced alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionic acid receptor endocytosis, a mechanism crucial for LTD expression. Thus, the BDNF pro-peptide is involved in synaptic plasticity that regulates a mechanism responsible for promoting LTD. The well-known BDNF polymorphism Val66Met affects human memory function. Here, the BDNF pro-peptide with Met mutation completely inhibits hippocampal LTD. These findings demonstrate functional roles for the BDNF pro-peptide and a naturally occurring human BDNF polymorphism in hippocampal synaptic depression.